

1. Population

| Population | 0 to <1 | | Total <1 | 1 to 4 | | Total 1 to 4 | 5 to 14 | | Total 5 to 14 | 15 to 49 | | Total 15 to 49 | ≥ 50 | | Total ≥ 50 | Grand total |
|--------------------|---------|----|----------|--------|-----|--------------|---------|-----|---------------|----------|-----|----------------|------|-----|------------|-------------|
| | M | F | | M | F | | M | F | | M | F | | M | F | | |
| Refugee population | 63 | 56 | 119 | 250 | 223 | 473 | 452 | 395 | 847 | 685 | 665 | 1,350 | 196 | 199 | 395 | 3,184 |
| % by age group | | | 4% | | | 15% | | | 27% | | | 42% | | | 12% | 100% |

* estimates from registered population

2. Mortality

2.1 Mortality

| Record ONE cause of mortality per case | 0 to 4 | | Total <5 | U5 % mort | 5 to 14 | | Total 5 to 14 | 15 to 49 | | Total 15 to 49 | ≥ 50 | | Total > 50 | Grand total | Crude % mort |
|---|----------|----------|----------|-------------|----------|----------|---------------|----------|----------|----------------|----------|----------|------------|-------------|--------------|
| | M | F | | | M | F | | M | F | | M | F | | | |
| 1. Lower Respiratory Tract Infection (LRTI) | 0 | 0 | 0 | | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | |
| 2. Watery diarrhoea | 0 | 0 | 0 | | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | |
| 3. Bloody diarrhoea | 0 | 0 | 0 | | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | |
| 4. Cardiovascular disorders | 0 | 0 | 0 | | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | |
| 5. Ischaemic heart disease | 0 | 0 | 0 | | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | |
| 6. Injury | 0 | 0 | 0 | | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | |
| 7. Cancer | 0 | 0 | 0 | | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | |
| 8. Chronic obstructive pulmonary disease (COPD) | 0 | 0 | 0 | | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | |
| 9. Tuberculosis (confirmed) | 0 | 0 | 0 | | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | |
| 10. Neonatal death | 0 | 0 | 0 | | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | |
| Other | 0 | 0 | 0 | | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | |
| Total | 0 | 0 | 0 | 100% | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 100% |
| % by age group | | | | | | | | | | | | | | | |

| Mortality indicators | Male | Female | Total |
|--------------------------------|------|--------|-------|
| Crude Mortality Rate (CMR) | 0.0 | 0.0 | 0.0 |
| Under 5 Mortality Rate (USMR) | 0.0 | 0.0 | 0.0 |
| Neonatal Mortality Rate (NNMR) | | | 0.0 |

2.2 Maternal death

| | |
|--|---|
| Number of maternal deaths reported | 0 |
| Number of maternal deaths investigated | 0 |
| Proportion of maternal deaths investigated | |

3. Morbidity

3.1 Staffing and service availability

| | |
|---|------|
| Number of full-time clinicians | 2 |
| Average number of full days on which each health facility was | 7.0 |
| Total number of outpatient consultations | 774 |
| Health facility utilisation rate | 12.6 |
| Consultations per clinician per day | 55 |

3.2 Acute health conditions

| Record NEW cases of disease only | 0 to 4 | | Total <5 | U5 Incid** | 5 to 14 | | Total 5 to 14 | 15 to 49 | | Total 15 to 49 | ≥ 50 | | Total ≥ 50 | Grand total | % by cause | Crude Incid** |
|--|-----------|-----------|------------|--------------|-----------|-----------|---------------|------------|------------|----------------|-----------|-----------|------------|-------------|-------------|---------------|
| | M | F | | | M | F | | M | F | | M | F | | | | |
| Upper Respiratory Tract Infection (URTI) | 55 | 49 | 104 | 175.7 | 62 | 52 | 114 | 57 | 61 | 118 | 8 | 7 | 15 | 351 | 49% | 110.2 |
| Lower Respiratory Tract Infection (LRTI) | 3 | 5 | 8 | 13.5 | 1 | 0 | 1 | 5 | 4 | 9 | 0 | 0 | 0 | 18 | 3% | 5.7 |
| Watery diarrhoea* | 9 | 5 | 14 | 23.6 | 1 | 5 | 6 | 2 | 2 | 4 | 0 | 0 | 0 | 24 | 3% | 7.5 |
| Bloody diarrhoea* | 1 | 0 | 1 | 1.7 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 1 | 0% | 0.3 |
| Malaria (suspected)* | 0 | 0 | 0 | 0.0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0% | 0.0 |
| Skin infection | 11 | 2 | 13 | 22.0 | 1 | 2 | 3 | 10 | 10 | 20 | 1 | 1 | 2 | 38 | 5% | 11.9 |
| Eye infection | 4 | 4 | 8 | 13.5 | 5 | 1 | 6 | 7 | 2 | 9 | 0 | 2 | 2 | 25 | 3% | 7.9 |
| Ear infection | 1 | 1 | 2 | 3.4 | 3 | 0 | 3 | 0 | 0 | 0 | 1 | 0 | 1 | 6 | 1% | 1.9 |
| Dental Conditions | 0 | 0 | 0 | 0.0 | 2 | 6 | 8 | 17 | 20 | 37 | 1 | 1 | 2 | 47 | 7% | 14.8 |
| Intestinal Worms | 0 | 0 | 0 | 0.0 | 2 | 0 | 2 | 0 | 0 | 0 | 0 | 0 | 0 | 2 | 0% | 0.6 |
| Acute Jaundice Syndrome* | 0 | 0 | 0 | 0.0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0% | 0.0 |
| Measles (suspected)* | 0 | 0 | 0 | 0.0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0% | 0.0 |
| Meningitis (suspected)* | 0 | 0 | 0 | 0.0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0% | 0.0 |
| Acute Flaccid Paralysis* | 0 | 0 | 0 | 0.0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0% | 0.0 |
| Tuberculosis (suspected) | 0 | 0 | 0 | 0.0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0% | 0.0 |
| Fever of unknown origin | 1 | 1 | 2 | 3.4 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 2 | 0% | 0.6 |
| Sexually Transmitted Infection (STI) | 0 | 0 | 0 | 0.0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0% | 0.0 |
| Urinary tract infection (UTI) | 0 | 0 | 0 | 0.0 | 0 | 1 | 1 | 6 | 7 | 13 | 1 | 0 | 1 | 15 | 2% | 4.7 |
| HIV/AIDS | 0 | 0 | 0 | 0.0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0% | 0.0 |
| Acute malnutrition | 0 | 0 | 0 | 0.0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0% | 0.0 |
| Leishmaniasis | 0 | 0 | 0 | 0.0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0% | 0.0 |
| Other | 12 | 9 | 21 | 35.5 | 7 | 7 | 14 | 55 | 90 | 145 | 7 | 4 | 11 | 191 | 27% | 60.0 |
| Total | 97 | 76 | 173 | 292.2 | 84 | 74 | 158 | 159 | 196 | 355 | 19 | 15 | 34 | 720 | 100% | 226.1 |
| % by age group | | | 24% | | | | 22% | | | 49% | | | 5% | 100% | | |

* indicates disease with epidemic potential. See guidance for alert thresholds and advice on how these should be interpreted. **incidence is cases per 1000 per week

3.3 Chronic health conditions

| Record NEW and REVISITS | 0 to 4 | | Total < 5 | 5 to 14 | | Total 5 to 14 | 15 to 49 | | Total 15 to 49 | ≥ 50 | | Total ≥ 50 | Grand total | % by cause | |
|--|----------|----------|-----------|----------|----------|---------------|----------|----------|----------------|----------|-----------|------------|-------------|-------------|--|
| | M | F | | M | F | | M | F | | M | F | | | | |
| Diabetes | 0 | 0 | 0 | 0 | 0 | 0 | 1 | 2 | 3 | 2 | 7 | 9 | 12 | 39% | |
| Hypertension | 0 | 0 | 0 | 0 | 0 | 0 | 2 | 1 | 3 | 4 | 9 | 13 | 16 | 52% | |
| Asthma | 0 | 0 | 0 | 0 | 0 | 0 | 1 | 0 | 1 | 0 | 0 | 0 | 1 | 3% | |
| Ischaemic heart disease | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0% | |
| Cardiovascular disease (other) | 0 | 0 | 0 | 0 | 0 | 0 | 1 | 0 | 1 | 0 | 0 | 0 | 1 | 3% | |
| Chronic obstructive pulmonary disease (COPD) | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0% | |
| Liver disease | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0% | |
| Thyroid problems | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0% | |
| Cancer | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0% | |
| Other | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 1 | 1 | 1 | 3% | |
| Total | 0 | 0 | 0 | 0 | 0 | 0 | 5 | 3 | 8 | 6 | 17 | 23 | 31 | 100% | |
| % by age group | | | 0% | | | 0% | | | 26% | | | 74% | 100% | | |

3.4 Mental health

| | 0 to 4 | | | 5 to 14 | | | 15 to 49 | | | ≥ 50 | | | Total ≥ 50 | Grand total | % by cause |
|---|----------|----------|--------------|----------|----------|------------------|----------|----------|-------------------|----------|----------|---------------|---------------|----------------|---------------|
| | M | F | Total < 5 | M | F | Total 5 to 14 | M | F | Total 15 to 49 | M | F | Total ≥ 50 | | | |
| 1. Epilepsy / seizures | 0 | 0 | 0 | 0 | 0 | 0 | 1 | 1 | 2 | 0 | 0 | 0 | 2 | 2 | 100% |
| 2. Alcohol and other substance abuse disorder | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0% |
| 3. Psychotic disorder | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0% |
| 4. Mental retardation / intellectual disability | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0% |
| 5. Severe emotional disorder | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0% |
| 6. Medically unexplained somatic complaint | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0% |
| Other | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0% |
| Total | 0 | 0 | 0 | 0 | 0 | 0 | 1 | 1 | 2 | 0 | 0 | 0 | 2 | 2 | 100% |
| % by age group | | | 0% | | | 0% | | | 100% | | | | 0% | 100% | |

3.5 Injuries

| | 0 to 4 | | | 5 to 14 | | | 15 to 49 | | | ≥ 50 | | | Total ≥ 50 | Grand total | % by cause |
|---------------------------|----------|----------|--------------|----------|----------|------------------|----------|----------|-------------------|----------|----------|---------------|---------------|----------------|---------------|
| | M | F | Total < 5 | M | F | Total 5 to 14 | M | F | Total 15 to 49 | M | F | Total ≥ 50 | | | |
| 1. Injury (intentional) | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0% |
| 2. Injury (unintentional) | 2 | 0 | 2 | 1 | 1 | 2 | 7 | 4 | 11 | 0 | 0 | 0 | 15 | 15 | 71% |
| 3. Burns | 0 | 1 | 1 | 0 | 1 | 1 | 2 | 2 | 4 | 0 | 0 | 0 | 6 | 6 | 29% |
| 4. Accident | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0% |
| Other | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0% |
| Total | 2 | 1 | 3 | 1 | 2 | 3 | 9 | 6 | 15 | 0 | 0 | 0 | 21 | 21 | 100% |
| % by age group | | | 14% | | | 14% | | | 71% | | | | 0% | 100% | |

4. Referrals

4.1 Referrals - general

| | 0 to 4 | 5 to 14 | 15 to 49 | ≥ 50 | Grand total |
|----------------|----------|----------|-----------|----------|----------------|
| Emergency | 0 | 1 | 4 | 0 | 5 |
| Elective | 4 | 6 | 18 | 0 | 28 |
| Other | 0 | 0 | 0 | 0 | 0 |
| Total | 4 | 7 | 22 | 0 | 33 |
| % by age group | 12% | 21% | 67% | 0% | 100% |

ND: no data submitted

5. Laboratory & other investigations

| | Total |
|-------------------------|----------|
| Haematology | 0 |
| Imaging (X-ray, MRI/CT) | 0 |
| Other | 0 |
| Total | 0 |

6. EPI and Vitamin A

| | <1 | ≥1 to <5 | Total |
|--|----|----------|-------|
| BCG | 0 | 0 | 0 |
| Hepatitis B birth dose | 0 | 0 | 0 |
| DPT I/HiB I/Hep B I* (Pentavalent I) | 0 | 0 | 0 |
| DPT II/HiB II* (Tetravalent) | 0 | 0 | 0 |
| DPT III/HiB III/Hep B II* (Pentavalent II) | 0 | 0 | 0 |
| Polio 0 | 0 | 0 | 0 |
| Polio I | 0 | 0 | 0 |
| Polio II | 0 | 0 | 0 |
| Polio III | 0 | 0 | 0 |
| Measles | 0 | 0 | 0 |
| MMR | 0 | 0 | 0 |
| Rotavirus 1 | 0 | 0 | 0 |
| Rotavirus 2 | 0 | 0 | 0 |
| Rotavirus 3 | 0 | 0 | 0 |
| Vitamin A | 0 | 0 | 0 |
| Additional vaccines | 0 | 0 | 0 |

| EPI Indicators (since December 2013) | | |
|--------------------------------------|------------------------|------|
| Total vaccinated this week | | 0 |
| a. Program dropout rate* | | -73% |
| b. Immunisation coverage | | |
| | BCG | 52% |
| | Hepatitis B birth dose | 48% |
| | DPT I | 41% |
| | DPT III | 71% |
| | Polio I | 34% |
| | Polio III | 71% |
| | Measles | 26% |
| | MMR** | 78% |
| | Rotavirus I | 15% |
| | Rotavirus III | 52% |
| c. Vitamin A coverage | | 58% |

*If program dropout <0%, please review data recording and reporting procedures for multi-dose vaccines
Targets: Program dropout, 0 to <10%. Immunisation coverage, ≥90%

**For most indicators, coverage estimates calculated for <1 years. MMR calculated for 1 to <2 years.

Note: If actual population living in the camps is lower than reported population, coverage indicators will be artificially depressed.

7. Reproductive Health

7.1 Antenatal care

| | < 18 | ≥ 18 | Total |
|--|----------------|------|-------|
| First antenatal visit < 1st trimester | | | |
| First antenatal visit > 1st trimester | | | |
| Number of pregnant women at time of delivery who received: | NO DATA | | |
| ≥ 4 ANC visits | | | |
| 2 TT doses | | | |
| anaemia screening | | | |

ND: no data submitted

| ANC Indicators | |
|--|--|
| Proportion of first time antenatal visits made < 1st trimester | |

7.2 Postnatal care

| | < 18 | ≥ 18 | Total |
|--|------|------|----------------|
| Women who attended 3 postnatal visits within 6 wks of delivery | | | NO DATA |

ND: no data submitted

7.3 Sexual and gender-based violence

| | < 18 | | ≥ 18 | | Total |
|--|------|---|------|---|-------|
| | M | F | M | F | |
| Total no rape survivors seen | 0 | 0 | 0 | 0 | 0 |
| No. of rape survivors examined within 72 hours | 0 | 0 | 0 | 0 | 0 |
| No. of rape survivors examined within 72 -120 hours | 0 | 0 | 0 | 0 | 0 |
| No. of rape survivors examined within 120 hrs - 2 weeks | 0 | 0 | 0 | 0 | 0 |
| No. of rape survivors examined after 2 weeks | 0 | 0 | 0 | 0 | 0 |
| No. of rape survivors given PEP within 72 hours | 0 | 0 | 0 | 0 | 0 |
| No. of female rape survivors given ECP within 120 hours | | | | | 0 |
| No. of rape survivors given STI presumptive treatment within 2 weeks | 0 | 0 | 0 | 0 | 0 |
| No. cases of trauma in health centre due to domestic violence | 0 | 0 | 0 | 0 | 0 |

ECP = Emergency Contraceptive Pill; PEP = Post-exposure Prophylaxis; STI = Sexually Transmitted Infection

| SGBV Indicators | | |
|--|--|-----|
| Incidence of reported rape | | 0.0 |
| Proportion of rape survivors who receive PEP < 72 hrs | | |
| Proportion of female rape survivors who receive ECP < 120 hrs | | |
| Proportion of rape survivors who receive STI presumptive treatment < 2 wks | | |

Data Sources

1. Health Information System

Publication date:

15 April 2014

Contact Information

Erbil
Dr Chimmentya Akrawi, DOH Erbil

Email: chimmentya@yahoo.com

Erbil
Sandra Harlass, UNHCR

Email: HARLASS@unhcr.org

We gratefully acknowledge the contributions of the following agencies:

