

1. Population

| Population | 0 to < 1 | | Total <1 | 1 to 4 | | Total 1 to 4 | 5 to 17 | | Total 5 to 17 | 18 to 59 | | Total 18 to 59 | > 60 | | Total > 60 | Grand total |
|--------------------|----------|-------|----------|--------|-------|--------------|---------|--------|---------------|----------|--------|----------------|------|-------|------------|-------------|
| | M | F | | M | F | | M | F | | M | F | | M | F | | |
| Refugee population | 1,660 | 1,581 | 3,240 | 6,119 | 5,774 | 11,894 | 14,687 | 13,914 | 28,601 | 16,438 | 13,724 | 30,162 | 890 | 1,248 | 2,138 | 76,035 |
| % by age group | | | 4% | | | 16% | | | 38% | | | 40% | | | 3% | 100% |

2. Mortality

2.1 Mortality

| Record ONE cause of mortality per case | 0 to 4 | | Total <5 | US % mort | 5 to 17 | | Total 5 to 17 | 18 to 59 | | Total 18 to 59 | > 60 | | Total > 60 | Grand total | Crude % mort |
|--|----------|-----------|-----------|-------------|----------|----------|---------------|----------|----------|----------------|-----------|-----------|------------|-------------|--------------|
| | M | F | | | M | F | | M | F | | M | F | | | |
| 1. Lower Respiratory Tract Infection (LRTI) | 0 | 2 | 2 | 11% | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 1 | 1 | 3 | 5% |
| 2. Watery diarrhoea | 0 | 0 | 0 | 0% | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0% |
| 3. Bloody diarrhoea | 0 | 0 | 0 | 0% | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0% |
| 4. Ischaemic heart disease | 0 | 0 | 0 | 0% | 0 | 0 | 0 | 1 | 0 | 1 | 3 | 1 | 4 | 5 | 8% |
| 5. Cardiovascular disorders (other) | 1 | 1 | 2 | 11% | 0 | 0 | 0 | 0 | 1 | 1 | 7 | 4 | 11 | 14 | 21% |
| 6. Cerebrovascular disease | 0 | 0 | 0 | 0% | 0 | 0 | 0 | 0 | 0 | 0 | 4 | 0 | 4 | 4 | 6% |
| 7. Chronic kidney disease | 0 | 0 | 0 | 0% | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0% |
| 8. Injury | 0 | 0 | 0 | 0% | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0% |
| 9. Cancer | 0 | 0 | 0 | 0% | 0 | 0 | 0 | 1 | 1 | 2 | 2 | 1 | 3 | 5 | 8% |
| 10. Chronic obstructive pulmonary disease (COPD) | 0 | 0 | 0 | 0% | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0% |
| 11. Tuberculosis (confirmed) | 0 | 0 | 0 | 0% | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0% |
| 12. Neonatal death | 3 | 5 | 8 | 44% | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 8 | 12% |
| 13. Other | 2 | 4 | 6 | 33% | 3 | 1 | 4 | 1 | 3 | 4 | 5 | 8 | 13 | 27 | 41% |
| Total | 6 | 12 | 18 | 100% | 3 | 1 | 4 | 3 | 5 | 8 | 21 | 15 | 36 | 66 | 100% |
| % by age group | | | 27% | | | | 6% | | | 12% | | | 55% | 100% | |

| Mortality indicators | Male | Female | Total |
|---|------|--------|-------|
| Crude Mortality Rate (CMR) (/1000/month) | 0.3 | 0.3 | 0.3 |
| Under 5 Mortality Rate (USMR) (/1000/month) | 0.3 | 0.5 | 0.4 |
| Neonatal Mortality Rate (NNMR) (/1000 livebirths) | | | 9.9 |

2.2 Maternal death

| | |
|--|---|
| Number of maternal deaths reported | 1 |
| Number of maternal deaths investigated | 0 |
| Proportion of maternal deaths investigated | |

3. Morbidity

3.1 Staffing and service availability

| | |
|---|---------|
| Number of full-time clinicians | 62.5 |
| Average number of full days on which each health facility was functioning | 6.2 |
| Total number of outpatient consultations | 181,762 |
| Health facility utilisation rate | 9.6 |
| Consultations per clinician per day | 36 |

3.2 Acute health conditions

| Record NEW cases of disease only | 0 to 4 | | Total <5 | US Incid | 5 to 17 | | Total 5 to 17 | 18 to 59 | | Total 18 to 59 | > 60 | | Total > 60 | Grand total | % by cause | Crude Incide |
|--|---------------|---------------|---------------|---------------|---------------|---------------|---------------|---------------|---------------|----------------|--------------|--------------|--------------|---------------|-------------|---------------|
| | M | F | | | M | F | | M | F | | M | F | | | | |
| Upper Respiratory Tract Infection (URTI) | 5,046 | 4,673 | 9,719 | 642.2 | 5,000 | 5,054 | 10,054 | 4,618 | 5,582 | 10,200 | 754 | 884 | 1,638 | 31,611 | 32% | 415.7 |
| Influenza-like illness (ILI) | 2,063 | 2,095 | 4,158 | 274.7 | 1,961 | 1,779 | 3,740 | 1,435 | 1,894 | 3,329 | 211 | 250 | 461 | 11,688 | 12% | 153.7 |
| Lower Respiratory Tract Infection (LRTI) | 1,146 | 792 | 1,938 | 128.1 | 735 | 767 | 1,502 | 692 | 764 | 1,456 | 236 | 236 | 472 | 5,368 | 6% | 70.6 |
| Watery diarrhoea* | 59 | 40 | 99 | 6.5 | 53 | 51 | 104 | 41 | 32 | 73 | 14 | 13 | 27 | 303 | 0% | 4.0 |
| Bloody diarrhoea* | 3 | 10 | 13 | 0.9 | 15 | 10 | 25 | 0 | 0 | 0 | 0 | 0 | 0 | 38 | 0% | 0.5 |
| Skin infection | 556 | 541 | 1,097 | 72.5 | 495 | 598 | 1,093 | 661 | 731 | 1,392 | 183 | 166 | 349 | 3,931 | 4% | 51.7 |
| Eye infection | 386 | 296 | 682 | 45.1 | 380 | 323 | 703 | 581 | 797 | 1,378 | 216 | 197 | 413 | 3,176 | 3% | 41.8 |
| Ear infection | 460 | 544 | 1,004 | 66.3 | 495 | 589 | 1,084 | 775 | 896 | 1,671 | 229 | 164 | 393 | 4,152 | 4% | 54.6 |
| Dental Conditions | 392 | 376 | 768 | 50.7 | 1,243 | 1,216 | 2,459 | 2,451 | 3,010 | 5,461 | 315 | 336 | 651 | 9,339 | 10% | 122.8 |
| Intestinal Worms | 262 | 284 | 546 | 36.1 | 303 | 305 | 608 | 96 | 93 | 189 | 6 | 1 | 7 | 1,350 | 1% | 17.8 |
| Acute Jaundice Syndrome* | 0 | 0 | 0 | 0.0 | 3 | 2 | 5 | 0 | 0 | 0 | 1 | 1 | 2 | 7 | 0% | 0.1 |
| Measles (suspected)* | 0 | 0 | 0 | 0.0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0% | 0.0 |
| Meningitis (suspected)* | 2 | 1 | 3 | 0.2 | 4 | 1 | 5 | 0 | 0 | 0 | 0 | 0 | 0 | 8 | 0% | 0.1 |
| Acute Flaccid Paralysis* | 0 | 0 | 0 | 0.0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0% | 0.0 |
| Leishmaniasis | 0 | 0 | 0 | 0.0 | 0 | 2 | 2 | 0 | 0 | 0 | 0 | 0 | 0 | 2 | 0% | 0.0 |
| Fever of unknown origin | 20 | 7 | 27 | 1.8 | 3 | 0 | 3 | 0 | 0 | 0 | 0 | 0 | 0 | 30 | 0% | 0.4 |
| Tuberculosis (suspected) | 0 | 0 | 0 | 0.0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0% | 0.0 |
| Other | 1,613 | 1,581 | 3,194 | 211.0 | 2,753 | 2,744 | 5,497 | 6,271 | 8,555 | 14,826 | 1,391 | 1,611 | 3,002 | 26,519 | 27% | 348.8 |
| Total | 12,008 | 11,240 | 23,248 | 1536.1 | 13,443 | 13,441 | 26,884 | 17,621 | 22,354 | 39,975 | 3,556 | 3,859 | 7,415 | 97,522 | 100% | 1282.6 |
| % by age group | | | 24% | | | | 28% | | | 41% | | | 8% | 100% | | |

* indicates disease with epidemic potential. See guidance for alert thresholds and advice on how these should be interpreted.

3.3 Chronic health conditions

| Record NEW and REVISITS | 0 to 4 | | Total < 5 | 5 to 17 | | Total 5 to 17 | 18 to 59 | | Total 18 to 59 | > 60 | | Total > 60 | Grand total | % by cause | |
|--|------------|------------|--------------|------------|------------|---------------|--------------|--------------|----------------|--------------|--------------|---------------|---------------|-------------|--|
| | M | F | | M | F | | M | F | | M | F | | | | |
| Diabetes | 1 | 1 | 2 | 47 | 57 | 104 | 937 | 1,349 | 2,286 | 948 | 1,120 | 2,068 | 4,460 | 16% | |
| Hypertension | 0 | 0 | 0 | 1 | 6 | 7 | 1,288 | 1,639 | 2,927 | 1,273 | 1,529 | 2,802 | 5,736 | 21% | |
| Asthma | 343 | 363 | 706 | 255 | 196 | 451 | 491 | 591 | 1,082 | 448 | 444 | 892 | 3,131 | 11% | |
| Ischaemic heart disease | 0 | 0 | 0 | 0 | 7 | 7 | 260 | 307 | 567 | 341 | 326 | 667 | 1,241 | 5% | |
| Cardiovascular disease (other) | 11 | 14 | 25 | 23 | 26 | 49 | 307 | 402 | 709 | 291 | 288 | 579 | 1,262 | 5% | |
| Chronic obstructive pulmonary disease (COPD) | 0 | 5 | 5 | 0 | 5 | 5 | 351 | 289 | 640 | 357 | 350 | 707 | 1,357 | 5% | |
| Liver disease | 1 | 0 | 1 | 0 | 0 | 0 | 7 | 19 | 26 | 9 | 4 | 13 | 40 | 0% | |
| Thyroid problems | 1 | 0 | 1 | 23 | 65 | 88 | 424 | 880 | 1,304 | 319 | 492 | 811 | 2,204 | 8% | |
| Cancer | 0 | 0 | 0 | 0 | 3 | 3 | 5 | 2 | 7 | 1 | 2 | 3 | 13 | 0% | |
| Other | 187 | 178 | 365 | 311 | 267 | 578 | 2,000 | 3,209 | 5,209 | 685 | 977 | 1,662 | 7,814 | 29% | |
| Total | 544 | 561 | 1,105 | 660 | 632 | 1,292 | 6,070 | 8,687 | 14,757 | 4,672 | 5,532 | 10,204 | 27,358 | 100% | |
| % by age group | | | 4% | | | 5% | | | 54% | | | 37% | 100% | | |

5.2 Referrals by service

| | | 0 to 4 | Total < 5 | 5 to 17 | Total 5 to 17 | 18 to 59 | Total 18 to 59 | > 60 | Total | Grand total |
|----|-------------------|--------|-----------|---------|---------------|----------|----------------|------|-------|-------------|
| 1. | Paediatrics | 131 | 131 | 0 | 0 | 0 | 0 | 0 | 0 | 131 |
| 2. | Internal medicine | 45 | 45 | 81 | 81 | 412 | 412 | 92 | 92 | 630 |
| 3. | Obstetric | 0 | 0 | 37 | 37 | 304 | 304 | 0 | 0 | 341 |
| 4. | Gynaecological | 0 | 0 | 37 | 37 | 93 | 93 | 1 | 1 | 131 |
| 5. | Surgical | 61 | 61 | 54 | 54 | 130 | 130 | 63 | 63 | 308 |
| 6. | Other | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 |

6. Laboratory

| | Total |
|--------------------------------------|--------|
| Number of laboratory tests conducted | 39,045 |

8. Reproductive Health

8.1 Antenatal care

| | < 18 | ≥ 18 | Total |
|---------------------------------------|------|------|-------|
| First antenatal visit < 1st trimester | 390 | 992 | 1,382 |
| First antenatal visit > 1st trimester | 76 | 670 | 746 |

ANC Indicators

| | |
|--|-----|
| Proportion of first time antenatal visits made < 1st trimester | 65% |
|--|-----|

8.2 Antenatal care in facility

| | | | | |
|--|-------------------|----|-----|-----|
| Number of pregnant women at time of delivery who received: | ≥ 4 ANC visits | 67 | 520 | 587 |
| | 2 TT doses | 66 | 507 | 573 |
| | anaemia screening | 74 | 682 | 756 |

| | |
|--|-----|
| Coverage of complete antenatal care | 73% |
| Coverage of antenatal tetanus immunisation | 71% |
| Coverage of anaemia screening in pregnancy | 94% |

8.3 Delivery care

| | < 18 | ≥ 18 | Total |
|---|------|------|-------|
| Live births | 75 | 732 | 807 |
| Still births | 1 | 6 | 7 |
| Low birth weight (<2500 g) | 0 | 3 | 3 |
| Attended by skilled health worker | 75 | 732 | 807 |
| Number of obstetric complications treated | 28 | 95 | 123 |
| Number of caesarean sections performed | 28 | 307 | 335 |

Delivery Indicators

| | |
|--|------|
| Crude birth rate (/1000 population/ month) | 3.5 |
| Proportion of births attended by skilled health worker (excl. TBA) | 100% |
| Proportion of low birth weight deliveries | 0% |
| Proportion of all births performed by caesarean section | 42% |
| Proportion of obstetric complications treated among girls < 18 | 23% |

8.4 Postnatal care

| | < 18 | ≥ 18 | Total |
|--|------|------|-------|
| At least three PNC visits within 6 weeks | 87 | 477 | 564 |

Postnatal Indicators

| | |
|-------------------------------------|-----|
| Coverage of complete postnatal care | 70% |
|-------------------------------------|-----|

8.5 Family planning

| | New | | Repeat | | Total |
|------------------------------|------|------|--------|-------|-------|
| | < 18 | ≥ 18 | < 18 | ≥ 18 | |
| Oral contraceptives | 82 | 524 | 542 | 1,058 | 2,206 |
| Emergency Contraceptive Pill | 0 | 0 | 0 | 0 | 0 |
| Injectable (Depo-Provera) | 2 | 43 | 47 | 102 | 194 |
| Implantable (Norplant) | 0 | 0 | 0 | 0 | 0 |
| Intra-Uterine Device (IUD) | 1 | 184 | 1 | 38 | 224 |
| Condom (Male) | 28 | 418 | 476 | 858 | 1,780 |
| Other | 10 | 19 | 0 | 17 | 46 |

| | < 18 | ≥ 18 |
|-----------------------------|------|-------|
| Family Planning Counseling* | 640 | 2,534 |

8.6 Sexual and gender-based violence

| | < 18 | | ≥ 18 | | Total |
|---|------|---|------|----|-------|
| | M | F | M | F | |
| Total no rape survivors seen | 0 | 0 | 0 | 0 | 0 |
| No. of rape survivors examined within 72 hours | 0 | 0 | 0 | 0 | 0 |
| No. of rape survivors examined within 72 -120 hours | 0 | 0 | 0 | 0 | 0 |
| No. of rape survivors examined within 120 hrs - 2 weeks | 0 | 0 | 0 | 0 | 0 |
| No. of rape survivors examined after 2 weeks | 0 | 0 | 0 | 0 | 0 |
| No of rape survivors given PEP within 72 hours | 0 | 0 | 0 | 0 | 0 |
| No of female rape survivors given ECP within 120 hours | | | | | 0 |
| No of rape survivors given STI presumptive treatment within 2 weeks | 0 | 0 | 0 | 0 | 0 |
| No. cases of trauma in health centre due to domestic violence | 0 | 2 | 0 | 17 | 19 |

ECP = Emergency Contraceptive Pill; PEP = Post-exposure Prophylaxis; STI = Sexually Transmitted Infection

SGBV Indicators

| | |
|--|-----|
| Incidence of reported rape (/10,000 population/year) | 0.0 |
| Proportion of rape survivors who receive PEP < 72 hrs | |
| Proportion of female rape survivors who receive ECP < 120 hrs | |
| Proportion of rape survivors who receive STI presumptive treatment < 2 wks | |

| | | | |
|---------------------|------------------------------|--------------------------|------------|
| Data Sources | 1. Health Information System | Publication date: | 07/06/2018 |
|---------------------|------------------------------|--------------------------|------------|

| | | |
|----------------------------|--|---|
| Contact Information | Za'atri Asaad Kadhum, UNHCR Email: kadhum@unhcr.org | Amman Adam Musa Khalifa, UNHCR Email: khalifaa@unhcr.org |
|----------------------------|--|---|

UNHCR gratefully acknowledges the support of the following partners:

