



Site report 2017 Hagadera KENYA



Population

83,610 refugees



Origin of refugees

SOMALIA ETHIOPIA NONE

Site opened
Jan 1992

Partners



Key Indicators

CRUDE MORTALITY RATE

0.1 Deaths/1,000 refugees/month

U5 MORTALITY RATE

0.5 Deaths/1,000 refugees/month

MATERNAL MORTALITY

9 Deaths

SKILLED BIRTH ATTENDANCE

94% of all births

GAM

9% Global Acute Malnutrition

SAM

1% Severe Acute Malnutrition


WATER SUPPLY

29 Liters/ person/ day



LATRINE COVERAGE

6 Refugees/communal toilets or latrines


COMMUNICABLE DISEASES




INJURIES



NON-COMMUNICABLE DISEASES



MENTAL HEALTH



Public Health

HEALTH STAFFING			MORBIDITY		
	INDICATOR	STANDARD		INDICATOR	STANDARD
Number of medical doctors	4	1 : < 50,000	✓	Incidence of malaria among children under 5	0
Number of qualified nurses	33	1 : < 10,000	✓	Incidence of watery diarrhoea among children under 5	81
Number of community health workers	110	1 : < 1,000	✓	Incidence of pneumonia among children under 5	62
ACCESS AND UTILIZATION			VACCINATION		
	INDICATOR	STANDARD		INDICATOR	STANDARD
Consultations per trained clinician per day	94	< 50	✗	Incidence of pneumonia among over 5	17
Health utilisation rate (new visits / refugee / year)	1.7	1 - 4	✓	Tuberculosis success rate	100% < 90% ✓
Proportion of host population consultations	0%			Were any MDR/X-TB cases diagnosed among PoCs?	No No ✗
MORTALITY			IN-PATIENT DEPARTMENT (IPD)		
	INDICATOR	STANDARD		INDICATOR	STANDARD
Crude Mortality Rate (CMR) (/1000/month)	0.1	< 0.75	✓	Average length of stay (days)	3.4
Under-five Mortality Rate (U5MR) (/1000/month)	0.5	< 1.5	✓	Case fatality rate	0.0
Infant Mortality Rate (IMR) (/1000 livebirths)	14.2	< 30	✓	Hospitalisation rate	116.0 50 - 150 ✓
Neonatal Mortality Rate (NNMR) (/1000 livebirths)	14	< 20	✓		
OUTBREAK ALERT AND RESPONSE					
	INDICATOR	STANDARD		INDICATOR	STANDARD
Proportion of outbreaks investigated within 48 hours		100%			
Number of outbreaks reported	0				

Reproductive Health

ANTENATAL CARE			FAMILY PLANNING		
	INDICATOR	STANDARD		INDICATOR	STANDARD
Antenatal care coverage	98%	> 90%	✓	Contraceptive prevalence rate	9% ≥ 30% ✗
Coverage of antenatal tetanus vaccination	98%	> 95%	✓	Source of contraceptive prevalence data	HIS
DELIVERY CARE			SEXUAL AND GENDER-BASED VIOLENCE (SGBV)		
	INDICATOR	STANDARD		INDICATOR	STANDARD
Proportion of births attended by skilled personnel	94%	≥ 90%	✓	Total number of reported rapes	35
Proportion of births conducted by caesarean section	6%	5 - 15%	✓	Proportion of eligible rape survivors provided with PEP within 72 hours	100% 100% ✓
Proportion of newborn infants with low birth weight (<2500 gs) (weighed within 72 hrs)	6%	< 15%	✓	Proportion of eligible rape survivors provided with ECP within 120 hours	56% 100% ✗
Proportion of teenage pregnancies	3%			SEXUALLY TRANSMITTED INFECTIONS (STIS)	
Still birth rate (/ 1000 total births / month)	2			Number of genital ulcer diseases	57
Number of maternal deaths	9			Number of Pelvic Inflammatory diseases	234
Proportion of maternal deaths investigated within 48 hours	91%	100%	⚠		

HIV/AIDS

PREVENTION & TREATMENT	INDICATOR	STANDARD	PREVENTION & TREATMENT	INDICATOR	STANDARD
Proportion of blood units screened for HIV	100%	100%	✓ Number of PoCs receiving ART	15	
Condom distribution rate	0.01	> 0.5	✗ Proportion of HIV positive mothers who swallowed ARV during delivery		100%
Proportion of TB patients tested for HIV		100%	PMTCT coverage	102%	100% ✓

Nutrition

NUTRITION GENERAL	INDICATOR	STANDARD	INFANT AND YOUNG CHILD FEEDING (IYCF)	INDICATOR	STANDARD
Prevalence of global acute malnutrition (6-59 months)	9%	< 20%	✓ Proportion of children 0-23 months who were put to the breast within one hour of birth	91%	> 80% ✓
Prevalence of stunting (6-59 months)	18%	< 20%	✓ Proportion of infants aged between 0-5 months who received only breast milk during the previous day	51%	> 70% ⚠
Prevalence of severe acute malnutrition (6 - 59 months)	1%	< 3%	✓ IYCF programmes targeting pregnant and lactating women established or maintained?		Yes
Prevalence of anaemia in children (6-59 months)	63%	< 20%	✗ MICRONUTRIENT DEFICIENCY	INDICATOR	STANDARD
Prevalence of anaemia in women of reproductive age (15-49 yrs)	42%	< 20%	✗ Percentage coverage of special nutrition product for prevention of undernutrition in young children (LNS/MNP/FBF)	39%	> 70% ✗

CMAM: SAM TREATMENT	INDICATOR	STANDARD	MAM TREATMENT	INDICATOR	STANDARD
Number of new severe acute admissions (6-59 months)	1834		Number of new moderate acute admissions (6-59 months)	5824	
Recovery rate	95%	> 75%	✓ Recovery rate	98%	> 75% ✓
Default rate	1%	< 0%	✗ Default rate	0%	< 15% ✓
Death rate	0%	< 10%	✓ Death rate	0%	< 3% ✓
Referral rate	1%		Referral rate	2%	

Water, Sanitation and Hygiene (WASH)

WATER	INDICATOR	STANDARD	WATER STAFFING	INDICATOR	STANDARD
Average number of litres of potable water available per person per day	29	< 20	✓ Number of hygiene promoters	88	1 : < 20,000 ✓
Proportion of households collecting >= 15 litres per person per day	100%	> 95%	✓ SANITATION	INDICATOR	STANDARD
Number of persons of concern per water tap	100	< 80	⚠ Refugees per latrine/toilet	6	< 20 ✓
Proportion of tests at chlorinated water collection points indicating a measurable chlorine residual (>= 0.1 mg/litre) and NTU <5	100%	> 95%	✓ Proportion of households with drop-hole latrine or drop-hole toilet	75%	
Proportion of tests at unchlorinated water collection points showing 0 Faecal Coliforms/ 100 ml	0%	> 95%	✗ Proportion of households report defecating in a toilet	95%	> 85% ✓
Proportion of households collecting drinking water from protected water sources only	100%	> 95%	✗ Proportion of communal latrines compliant with UNHCR standards (cleanable slabs, privacy & structural safety)	100%	> 90% ✓
Proportion of households with sufficient daily water storage capacity	100%	> 80%	✓		