

Highlights

- From 27 April to 6 July 2017, 291 554 suspected cholera cases and 1 678 deaths (CFR: 0.6%) have been reported in 95.6% (22/23) of Yemen governorates, and 86.5% (288/333) of the districts.
- For the first time since the beginning of this outbreak second wave, four cases have been reported in Say'on governorate. All were adults. Two of these cases were confirmed with rapid diagnostic tests.

Geographical distribution of cases

The four most affected governorates were Amanat Al Asimah, Al Hudaydah, Hajjah and Amran with 47.4% (138 279/291 554) of the cases reported **since 27 April 2017**. Amran, Al Dhale'e and Al Mahwit governorates had the highest attack rates (20.2‰, 19.4‰ and 19.2‰ respectively), and Raymah, Hajjah and Ibb governorates the highest case fatality ratios (1.4%, 1.0% and 1.0% respectively) (see table).

Number of suspected cholera cases & deaths, AR and CFR by governorate, Yemen, 27 April – 6 July 2017

Governorates	Cases	Deaths	CFR (%)	Attack Rate (‰)
Amanat Al Asimah	40 477	54	0.1	12.2
Al-Hudaydah	35 646	193	0.5	10.7
Hajjah	31 329	312	1.0	14.1
Amran	30 827	146	0.5	20.2
Sana'a	23 017	109	0.5	18.4
Ibb	22 737	223	1.0	7.4
Taizz	21 421	146	0.7	7.1
Dhamar	18 124	112	0.6	8.5
Al Dhale'e	14 628	65	0.4	19.4
Al Mahwit	14 572	108	0.7	19.2
Al Bayda	8 898	24	0.3	11.6
Aden	8 519	46	0.5	8.9
Abyan	7 293	27	0.4	11.9
Raymah	5 517	78	1.4	8.7
Lahj	4 063	16	0.4	3.9
Al_Jawf	2 208	13	0.6	3.4
Ma'areb	1 279	4	0.3	3.6
Sa'ada	466	1	0.2	0.5
AL Mahrah	338	1	0.3	2.1
Shabwah	189	-	-	0.3
Say'on	4	-	-	0.0
Moklla	2	-	-	0.0
Total	291 554	1 678	0.6	9.8

Al Hali (Al Hodaydah gov., 12 245 cases, 21 deaths), Bani Al Harith (Amanat Al Asimah gov., 8 004 cases, 10 deaths), Ma'ain (Amanat Al Asimah gov., 7 800, 7 deaths) and As Sabain (Amanat Al Asimah gov., 6 097 cases, 9 deaths) were still the four most affected districts.

For further information:

Dr Xavier de Radiguès
C: +967 738 445 522

Dr Ahmed Zouiten
zouitena@who.int

Note: This report is an update to the weekly epidemiology bulletin.