

Highlights

- From 27 April to 7 July 2017, 297 438 suspected cholera cases and 1 706 deaths (CFR: 0.6%) have been reported in 95.6% (22/23) of Yemen governorates, and 86.5% (288/333) of the districts.

Geographical distribution of cases

The four most affected governorates were Amanat Al Asimah, Al Hudaydah, Hajjah and Amran with 47.3% (140 750/297 438) of the cases reported **since 27 April 2017**. Amran, Al Dhaele'e and Al Mahwit governorates had the highest attack rates (20.5‰, 20.1‰ and 19.6‰ respectively), and Raymah, Hajjah and Ibb governorates the highest case fatality ratios (1.4%, 1.0% and 1.0% respectively) (see table).

Number of suspected cholera cases & deaths, AR and CFR by governorate, Yemen, 27 April – 7 July 2017

Governorates	Cases	Deaths	CFR (%)	Attack Rate (‰)
Amanat Al Asimah	40 840	54	0.1	12.3
Al-Hudaydah	36 379	196	0.5	10.9
Hajjah	31 907	325	1.0	14.4
Amran	31 424	147	0.5	20.5
Sana'a	23 261	110	0.5	18.6
Ibb	23 129	225	1.0	7.5
Taizz	21 836	146	0.7	7.2
Dhamar	18 840	113	0.6	8.9
Al Dhaele'e	15 166	70	0.5	20.1
Al Mahwit	14 921	108	0.7	19.6
Al Bayda	9 028	24	0.3	11.7
Aden	8 948	48	0.5	9.4
Abyan	7 582	27	0.4	12.4
Raymah	5 534	78	1.4	8.7
Lahj	4 072	16	0.4	3.9
Al_Jawf	2 209	13	0.6	3.4
Ma'areb	1 335	4	0.3	3.7
Sa'ada	477	1	0.2	0.5
AL Mahrah	355	1	0.3	2.2
Shabwah	189	-	-	0.3
Say'on	4	-	-	0.0
Moklla	2	-	-	0.0
Total	297 438	1 706	0.6	10.0

Al Hali (Al Hudaydah gov., 12 363 cases, 22 deaths), Bani Al Harith (Amanat Al Asimah gov., 8 095 cases, 10 deaths), Ma'ain (Amanat Al Asimah gov., 7 850, 7 deaths) and As Sabain (Amanat Al Asimah gov., 6 149 cases, 9 deaths) were still the four most affected districts.

For further information:

Dr Xavier de Radiguès
C: +967 738 445 522

Lauren O'Connor
Communication officer
occonnorl@who.int

Note: This report is an update to the weekly epidemiology bulletin.